

On quasi-Jacobi and Jacobi-quasi bialgebroids

J. M. Nunes da Costa
 Departamento de Matemática
 Universidade de Coimbra
 Apartado 3008
 3001-454 Coimbra - Portugal
 E-mail: jmcosta@mat.uc.pt

F. Petalidou
 Faculty of Sciences and Technology
 University of Peloponnese
 22100 Tripoli - Greece
 E-mail: petalido@uop.gr

Abstract

We study quasi-Jacobi and Jacobi-quasi bialgebroids and their relationships with twisted Jacobi and quasi Jacobi manifolds. We show that we can construct quasi-Lie bialgebroids from quasi-Jacobi bialgebroids, and conversely, and also that the structures induced on their base manifolds are related via a “quasi Poissonization”.

Keywords: Quasi-Jacobi bialgebroid, Jacobi-quasi bialgebroid, quasi-Jacobi bialgebra, twisted Jacobi manifold, quasi Jacobi manifold.

A.M.S. classification (2000): 17B62; 17B66; 53D10; 53D17.

1 Introduction

The notion of *quasi-Lie bialgebroid* was introduced in [15]. It is a structure on a pair (A, A^*) of vector bundles, in duality, over a differentiable manifold M that is defined by a Lie algebroid structure on A^* , a skew-symmetric bracket on the space of smooth sections of A and a bundle map $a : A \rightarrow TM$, satisfying some compatibility conditions. These conditions are expressed in terms of a section φ of $\bigwedge^3 A^*$, which turns out to be an obstruction to the Lie bialgebroid structure on (A, A^*) . A quasi-Lie bialgebroid will be denoted by (A, A^*, φ) . In the case where A is a Lie algebroid, its dual vector bundle A^* is equipped with a skew-symmetric bracket on its space of smooth sections and a bundle map $a_* : A^* \rightarrow TM$ and the compatibility conditions are expressed in terms of a section Q of $\bigwedge^3 A$, the triple (A, A^*, Q) is called a *Lie-quasi bialgebroid* [9]. When $\varphi = 0$ and $Q = 0$, quasi-Lie and Lie-quasi bialgebroids are just Lie bialgebroids. We note that, while the dual of a Lie bialgebroid is itself a Lie bialgebroid, the dual of a quasi-Lie bialgebroid is a Lie-quasi bialgebroid, and conversely [9]. The quasi-Lie and Lie-quasi bialgebroids are particular cases of proto-bialgebroids [9]. As in the case

of a Lie bialgebroid, the doubles $A \oplus A^*$ of a quasi-Lie and of a Lie-quasi bialgebroid are endowed with a Courant algebroid structure [15], [9].

It was shown in [16] that the theory of quasi-Lie bialgebroids is the natural framework in which we can treat *twisted Poisson manifolds*. These structures were introduced in [17], under the name of Poisson manifolds with a closed 3-form background, motivated by problems of string theory [14] and of topological field theory [8].

The notion of *Jacobi bialgebroid* and the equivalent one of *generalized Lie bialgebroid* were introduced, respectively, in [3] and [5], in such a way that a Jacobi bialgebroid is canonically associated to a Jacobi manifold and conversely. A Jacobi bialgebroid over M is a pair $((A, \phi), (A^*, W))$ of Lie algebroids over M , in duality, endowed with 1-cocycles $\phi \in \Gamma(A^*)$ and $W \in \Gamma(A)$ in their Lie algebroid cohomology complexes with trivial coefficients, respectively, that satisfy a compatibility condition. Also, its double $(A \oplus A^*, \phi + W)$ is endowed with a Courant-Jacobi algebroid structure [4], [11].

In order to adapt to the framework of Jacobi manifolds the concepts of twisted Poisson manifold and quasi-Lie bialgebroid, we have recently introduced in [12] the notions of *twisted Jacobi manifold* and *quasi-Jacobi bialgebroid*. The purpose of the present paper is to develop the theory of quasi-Jacobi bialgebroids, as well as of its dual concept of *Jacobi-quasi bialgebroids*, and to establish a very close relationship between quasi-Jacobi and quasi-Lie bialgebroids.

The paper contains four sections, besides the Introduction, and one Appendix (section 5). In section 2 we recall the definition of quasi-Jacobi bialgebroid, we present some basic results established in [12], we develop the examples of quasi-Jacobi and Jacobi-quasi bialgebroids associated to twisted Jacobi manifolds and to quasi Jacobi manifolds, and, finally, we study the triangular quasi-Jacobi bialgebroids. Section 3 is devoted to the study of the structures induced on the base manifolds of quasi-Jacobi and Jacobi-quasi bialgebroids. Several examples are presented. In section 4 we establish a one to one correspondence between quasi-Jacobi bialgebroid structures $((A, \phi), (A^*, W), \varphi)$ over a manifold M and quasi-Lie bialgebroid structures $(\tilde{A}, \tilde{A}^*, \tilde{\varphi})$ over $\tilde{M} = M \times \mathbb{R}$. Also, we prove that the structure induced on $\tilde{M} = M \times \mathbb{R}$ by $(\tilde{A}, \tilde{A}^*, \tilde{\varphi})$ is the “quasi Poissonization” of the structure induced on M by $((A, \phi), (A^*, W), \varphi)$. The dual version of these results is also presented. Finally, in the Appendix, we define the action of a Lie algebroid with 1-cocycle on a differentiable manifold, a concept that is used in the paper.

Notation: If (A, ϕ) is a Lie algebroid with 1-cocycle ϕ , we denote by d^ϕ the differential operator d of A modified by ϕ , i.e., $d^\phi \alpha = d\alpha + \phi \wedge \alpha$, for any $\alpha \in \Gamma(\bigwedge^k A^*)$. Moreover, we denote by δ the usual de Rham differential operator on a manifold M and by d the differential operator of the Lie algebroid $TM \times \mathbb{R}$, $d(\alpha, \beta) = (\delta\alpha, -\delta\beta)$, for $(\alpha, \beta) \in \Gamma(\bigwedge^k(T^*M \times \mathbb{R})) \equiv \Gamma(\bigwedge^k T^*M) \times \Gamma(\bigwedge^{k-1} T^*M)$. We also consider the identification $\Gamma(\bigwedge^k(TM \times \mathbb{R})) \equiv \Gamma(\bigwedge^k TM) \times \Gamma(\bigwedge^{k-1} TM)$.

2 Quasi-Jacobi and Jacobi-quasi bialgebroids

Let $((A, \phi), (A^*, W))$ be a pair of dual vector bundles over a differentiable manifold M endowed with a 1-form ϕ and W , respectively, and φ a 3-form of A .

Definition 2.1 A quasi-Jacobi bialgebroid structure on $((A, \phi), (A^*, W), \varphi)$ consists of a Lie algebroid structure with 1-cocycle $([\cdot, \cdot]_*, a_*, W)$ on A^* , a bundle map $a : A \rightarrow TM$ and a skew-symmetric operation $[\cdot, \cdot]$ on $\Gamma(A)$ satisfying, for all $X, Y, Z \in \Gamma(A)$ and $f \in C^\infty(M, \mathbb{R})$, the following conditions:

- 1) $[X, fY] = f[X, Y] + (a(X)f)Y$;
- 2) $a([X, Y]) = [a(X), a(Y)] - a_*(\varphi(X, Y, \cdot))$;
- 3) $[[X, Y], Z] + c.p. = -d_*^W(\varphi(X, Y, Z)) - ((i_{\varphi(X, Y, \cdot)}d_*^W Z) + c.p.)$;
- 4) $d\phi - \varphi(W, \cdot, \cdot) = 0$, where d is the quasi-differential operator on $\Gamma(\wedge A^*)$ determined by the structure $([\cdot, \cdot], a)$ on A ;
- 5) $d^\phi\phi = 0$, d^ϕ being the quasi-differential operator modified by ϕ ;
- 6) $d_*^W[P, Q]^\phi = [d_*^W P, Q]^\phi + (-1)^{p+1}[P, d_*^W Q]^\phi$, where $[\cdot, \cdot]^\phi$ is the bracket on $\Gamma(\wedge A)$ modified by ϕ , $P \in \Gamma(\wedge^p A)$ and $Q \in \Gamma(\wedge A)$.

As in the case of quasi-Lie and Lie-quasi bialgebroids, by interchanging the roles of (A, ϕ) and (A^*, W) in the above definition, we obtain the notion of *Jacobi-quasi bialgebroid* over a differentiable manifold M . We have: If $((A, \phi), (A^*, W), \varphi)$ is a quasi-Jacobi bialgebroid over a differentiable manifold M , then $((A^*, W), (A, \phi), \varphi)$ is a Jacobi-quasi bialgebroid over M , and conversely.

In the case where both 1-cocycles ϕ and W are zero, we recover, from Definition 2.1, the notion of quasi-Lie bialgebroid. On the other hand, if $\varphi = 0$, then $((A, \phi), (A^*, W), 0) \equiv ((A, \phi), (A^*, W))$ is a Jacobi bialgebroid over M .

Remark 2.2 In [12], we proved that the double of a quasi-Jacobi bialgebroid is a Courant-Jacobi algebroid ([4], [11]). By a similar computation, we may conclude that the double of a Jacobi-quasi bialgebroid is also a Courant-Jacobi algebroid.

The rest of this section is devoted to some important examples of quasi-Jacobi and Jacobi-quasi bialgebroids.

2.1 Quasi-Jacobi and Jacobi-quasi bialgebras

A *quasi-Jacobi bialgebra* is a quasi-Jacobi bialgebroid over a point, that is a triple $((\mathcal{G}, \phi), (\mathcal{G}^*, W), \varphi)$, where $(\mathcal{G}^*, [\cdot, \cdot]_*, W)$ is a real Lie algebra of finite dimension with 1-cocycle $W \in \mathcal{G}$ in its Chevalley-Eilenberg cohomology, $(\mathcal{G}, \phi)$ is the dual space of $\mathcal{G}^*$ endowed with a bilinear skew-symmetric bracket $[\cdot, \cdot]$ and an element $\phi \in \mathcal{G}^*$ and $\varphi \in \wedge^3 \mathcal{G}^*$, such that conditions 3)-6) of Definition 2.1 are satisfied.

By dualizing the above notion, we get a *Jacobi-quasi bialgebra*, i.e. a Jacobi-quasi bialgebroid over a point.

In the particular case where $\varphi = 0$, we recover the concept of Jacobi bialgebra [5]. When $\phi = 0$ and $W = 0$, we recover the notion of quasi-Lie bialgebra due to Drinfeld [2].

We postpone the study of quasi-Jacobi bialgebras to a future paper. We believe that they can be considered as the infinitesimal invariants of Lie groups endowed with a certain type of twisted Jacobi structures that can be constructed from the solutions of a twisted Yang-Baxter equation.

2.2 The quasi-Jacobi and the Jacobi-quasi bialgebroids of a twisted Jacobi manifold

We recall that a *twisted Jacobi manifold* [12] is a differentiable manifold M equipped with a section (Λ, E) of $\bigwedge^2(TM \times \mathbb{R})$ and a 2-form ω such that

$$\frac{1}{2}[(\Lambda, E), (\Lambda, E)]^{(0,1)} = (\Lambda, E)^\#(\delta\omega, \omega),^1 \quad (1)$$

where $[\cdot, \cdot]^{(0,1)}$ denotes the Schouten bracket of the Lie algebroid $(TM \times \mathbb{R}, [\cdot, \cdot], \pi)$ over M modified by the 1-cocycle $(0, 1)$ and $(\Lambda, E)^\#$ is the natural extension of the homomorphism of $C^\infty(M, \mathbb{R})$ -modules $(\Lambda, E)^\# : \Gamma(T^*M \times \mathbb{R}) \rightarrow \Gamma(TM \times \mathbb{R})$, $(\Lambda, E)^\#(\alpha, f) = (\Lambda^\#(\alpha) + fE, -\langle \alpha, E \rangle)$, to a homomorphism from $\Gamma(\bigwedge^k(T^*M \times \mathbb{R}))$ to $\Gamma(\bigwedge^k(TM \times \mathbb{R}))$, $k \in \mathbb{N}$, given, for any $(\eta, \xi) \in \Gamma(\bigwedge^k(T^*M \times \mathbb{R}))$ and $(\alpha_1, f_1), \dots, (\alpha_k, f_k) \in \Gamma(T^*M \times \mathbb{R})$, by

$$\begin{aligned} & (\Lambda, E)^\#(\eta, \xi)((\alpha_1, f_1), \dots, (\alpha_k, f_k)) \\ &= (-1)^k(\eta, \xi)((\Lambda, E)^\#(\alpha_1, f_1), \dots, (\Lambda, E)^\#(\alpha_k, f_k)) \end{aligned}$$

and, for all $f \in C^\infty(M, \mathbb{R})$, by $(\Lambda, E)^\#(f) = f$.

Examples of twisted Jacobi manifolds are *twisted exact Poisson manifolds* and *twisted locally conformal symplectic manifolds*, both of them presented in [12], and also *twisted contact Jacobi manifold* described in [13].

If $(M, (\Lambda, E), \omega)$ is a twisted Jacobi manifold and f a function on M that never vanishes, we can define a new twisted Jacobi structure $((\Lambda^f, E^f), \omega^f)$ on M , which is said to be *f-conformal* to $((\Lambda, E), \omega)$, by setting

$$\Lambda^f = f\Lambda; \quad E^f = \Lambda^\#(\delta f) + fE; \quad \omega^f = \frac{1}{f}\omega.$$

In the sequel, let $(M, (\Lambda, E), \omega)$ be a twisted Jacobi manifold and $(T^*M \times \mathbb{R}, [\cdot, \cdot]_{(\Lambda, E)}^\omega, \pi \circ (\Lambda, E)^\#, (-E, 0))$ its canonically associated Lie algebroid with 1-cocycle, [12]. The Lie bracket $[\cdot, \cdot]_{(\Lambda, E)}^\omega$ on $\Gamma(T^*M \times \mathbb{R})$ is given, for all $(\alpha, f), (\beta, g) \in \Gamma(T^*M \times \mathbb{R})$, by

$$\begin{aligned} [(\alpha, f), (\beta, g)]_{(\Lambda, E)}^\omega &= [(\alpha, f), (\beta, g)]_{(\Lambda, E)} \\ &\quad + (\delta\omega, \omega)((\Lambda, E)^\#(\alpha, f), (\Lambda, E)^\#(\beta, g), \cdot), \end{aligned}$$

¹Since, for any $(\varphi, \omega) \in \Gamma(\bigwedge^3(T^*M \times \mathbb{R}))$, $d^{(0,1)}(\varphi, \omega) = (\delta\varphi, \varphi - \delta\omega)$ and $d^{(0,1)}(\varphi, \omega) = (0, 0) \Leftrightarrow \varphi = \delta\omega$, equation (1) means that $\frac{1}{2}[(\Lambda, E), (\Lambda, E)]^{(0,1)}$ is the image by $(\Lambda, E)^\#$ of a $d^{(0,1)}$ -closed 3-form of $TM \times \mathbb{R}$.

where $[\cdot, \cdot]_{(\Lambda, E)}$ is the usual bracket on $\Gamma(T^*M \times \mathbb{R})$ associated to a section (Λ, E) of $\bigwedge^2(TM \times \mathbb{R})$ ([7], [5]):

$$\begin{aligned} [(\alpha, f), (\beta, g)]_{(\Lambda, E)} &= \mathcal{L}_{(\Lambda, E)^\#(\alpha, f)}^{(0,1)}(\beta, g) - \mathcal{L}_{(\Lambda, E)^\#(\beta, g)}^{(0,1)}(\alpha, f) \\ &\quad - d^{(0,1)}((\Lambda, E)((\alpha, f), (\beta, g))). \end{aligned} \quad (2)$$

We consider, on the vector bundle $TM \times \mathbb{R} \rightarrow \mathbb{R}$, the Lie algebroid structure over M with 1-cocycle $([\cdot, \cdot], \pi, (0, 1))$ and also a new bracket $[\cdot, \cdot]'$ on the space of its smooth sections given, for all $(X, f), (Y, g) \in \Gamma(TM \times \mathbb{R})$, by

$$[(X, f), (Y, g)]' = [(X, f), (Y, g)] - (\Lambda, E)^\#((\delta\omega, \omega)((X, f), (Y, g), \cdot)).$$

We have shown in [12] that the triple $((TM \times \mathbb{R}, [\cdot, \cdot]', \pi, (0, 1)), (T^*M \times \mathbb{R}, [\cdot, \cdot]_{(\Lambda, E)}^\omega, \pi \circ (\Lambda, E)^\#, (-E, 0)), (\delta\omega, \omega))$ is a quasi-Jacobi bialgebroid over M . Furthermore, we have:

Theorem 2.3 *The triple $((TM \times \mathbb{R}, [\cdot, \cdot]', \pi, (0, 1)), (T^*M \times \mathbb{R}, [\cdot, \cdot]_{(\Lambda, E)}^\omega, \pi \circ (\Lambda, E)^\#, (-E, 0)), (\Lambda, E)^\#(\delta\omega, \omega))$ is a Jacobi-quasi bialgebroid over M .*

Proof. It suffices to check that the dual versions of all conditions of Definition 2.1 are satisfied. Condition 1) can be checked directly, using (2). For 2), we take into account that $((\Lambda, E), \omega)$ is a twisted Jacobi structure, hence (1) holds, and we apply the general formula

$$\begin{aligned} (\Lambda, E)^\#([(\alpha, f), (\beta, g)]_{(\Lambda, E)}) &= [(\Lambda, E)^\#(\alpha, f), (\Lambda, E)^\#(\beta, g)] \\ &\quad - \frac{1}{2}[(\Lambda, E), (\Lambda, E)]^{(0,1)}((\alpha, f), (\beta, g), \cdot). \end{aligned} \quad (3)$$

By projection, we obtain 2). Condition 3) can be checked directly, after a long computation. In order to prove 4), we remark that the quasi-differential operator d_* determined by $([\cdot, \cdot]_{(\Lambda, E)}, \pi \circ (\Lambda, E)^\#)$ is given [5], for all $(R, S) \in \Gamma(\bigwedge^k(TM \times \mathbb{R}))$, by

$$d_*(R, S) = ([\Lambda, R] + kE \wedge R + \Lambda \wedge S, -[\Lambda, R] + (1 - k)E \wedge S + [E, R]).$$

So, $d_*(-E, 0) = ([E, \Lambda], 0)$, and since $(M, (\Lambda, E), \omega)$ is a twisted Jacobi manifold,

$$\begin{aligned} d_*(-E, 0) &= ([E, \Lambda], 0) = \frac{1}{2}[(\Lambda, E), (\Lambda, E)]^{(0,1)}((0, 1), \cdot, \cdot) \\ &= ((\Lambda, E)^\#(\delta\omega, \omega))((0, 1), \cdot, \cdot). \end{aligned}$$

On the other hand, since $d_*^{(-E, 0)}(R, S) = [(\Lambda, E), (R, S)]^{(0,1)}$, we have

$$\begin{aligned} d_*^{(-E, 0)}((\Lambda, E)^\#(\delta\omega, \omega)) &= [(\Lambda, E), (\Lambda, E)^\#(\delta\omega, \omega)]^{(0,1)} \\ &= \frac{1}{2}[(\Lambda, E), [(\Lambda, E), (\Lambda, E)]^{(0,1)}]^{(0,1)} = 0, \end{aligned}$$

whence we get condition 5). Finally, 6) can be established, as in the proof of Theorem 8.2 in [12], by a straightforward but long computation. $\square$

In the case of twisted Poisson manifolds, the previous results were treated in [16] and [9].

2.3 The Jacobi-quasi bialgebroid of a quasi Jacobi manifold

Let $(\mathcal{G}, [\cdot, \cdot])$ be a Lie algebra, ϕ a 1-cocycle in its Chevalley-Eilenberg cohomology and $(\cdot, \cdot)$ a nondegenerate symmetric bilinear invariant form on $\mathcal{G}$. We denote by ψ the canonical 3-form on $\mathcal{G}$ defined by $\psi(X, Y, Z) = \frac{1}{2}(X, [Y, Z])$, for all $X, Y, Z \in \mathcal{G}$, and by $Q_\psi \in \bigwedge^3 \mathcal{G}$ its dual trivector that is given, for all $\mu, \nu, \xi \in \mathcal{G}^*$, by

$$Q_\psi(\mu, \nu, \xi) = \psi(X_\mu, X_\nu, X_\xi),$$

where X_μ, X_ν, X_ξ are, respectively, dual to μ, ν, ξ via $(\cdot, \cdot)$.

A $(\mathcal{G}, \phi)$ -manifold M is a differentiable manifold on which $(\mathcal{G}, \phi)$ acts infinitesimally by $a^\phi : \mathcal{G} \rightarrow TM \times \mathbb{R}$, $a^\phi(X) = a(X) + \langle \phi, X \rangle$, for all $X \in \mathcal{G}$ (see Appendix). We keep the same notation a^ϕ for the induced maps on exterior algebras.

Let M be a $(\mathcal{G}, \phi)$ -manifold. A section $(\Lambda, E) \in \Gamma(\bigwedge^2(TM \times \mathbb{R}))$ is said to be *invariant* (under the infinitesimal action a^ϕ) if, for any $X \in \mathcal{G}$,

$$\mathcal{L}_{a^\phi(X)}^{(0,1)}(\Lambda, E) = (0, 0).$$

A natural generalization of the notion of quasi Poisson manifold, given in [1], is the concept of $(\mathcal{G}, \phi)$ -quasi Jacobi manifold, that we introduce as follows.

Definition 2.4 A $(\mathcal{G}, \phi)$ -quasi Jacobi manifold is a $(\mathcal{G}, \phi)$ -manifold M equipped with an invariant section $(\Lambda, E) \in \Gamma(\bigwedge^2(TM \times \mathbb{R}))$ such that

$$\frac{1}{2}[(\Lambda, E), (\Lambda, E)]^{(0,1)} = a^\phi(Q_\psi).$$

A long, but not difficult computation, leads us to the following:

Theorem 2.5 Let (M, Λ, E) be a $(\mathcal{G}, \phi)$ -quasi Jacobi manifold. Then, $((TM \times \mathbb{R}, [\cdot, \cdot], \pi, (0, 1)), (T^*M \times \mathbb{R}, [\cdot, \cdot]_{(\Lambda, E)}, \pi \circ (\Lambda, E)^\#, (-E, 0)), a^\phi(Q_\psi))$ is a Jacobi-quasi bialgebroid over M .

Remark 2.6 If M is a $\mathcal{G}$ -manifold equipped with a quasi Poisson structure, i.e an invariant bivector field Λ on M such that $[\Lambda, \Lambda] = 2a(Q_\psi)$, a similar result holds: The triple $((TM, [\cdot, \cdot], id), (T^*M, [\cdot, \cdot]_\Lambda, \Lambda^\#), a(Q_\psi))$ is a Lie-quasi bialgebroid over M , $[\cdot, \cdot]_\Lambda$ being the Koszul bracket associated to Λ .

2.4 Triangular quasi-Jacobi and Jacobi-quasi bialgebroids

Let $(A, [\cdot, \cdot], a, \phi)$ be a Lie algebroid with 1-cocycle over a differentiable manifold M , Π a section of $\bigwedge^2 A$ and Q a trivector on A such that

$$\frac{1}{2}[\Pi, \Pi]^\phi = Q. \quad (4)$$

We shall discuss what happens on the dual vector bundle A^* of A when we consider the vector bundle map $a_* : A^* \rightarrow TM$, $a_* = a \circ \Pi^\#$, $\Pi^\# : A^* \rightarrow A$ being the bundle map associated to Π , and the Koszul bracket $[\cdot, \cdot]_\Pi$ on the space $\Gamma(A^*)$ of its smooth sections given, for all $\alpha, \beta \in \Gamma(A^*)$, by

$$[\alpha, \beta]_\Pi = \mathcal{L}_{\Pi^\#(\alpha)}^\phi \beta - \mathcal{L}_{\Pi^\#(\beta)}^\phi \alpha - d^\phi(\Pi(\alpha, \beta)). \quad (5)$$

Let us set $W = -\Pi^\#(\phi)$. Taking into account that, for all $\alpha, \beta, \gamma \in \Gamma(A^*)$,

$$[[\alpha, \beta]_\Pi, \gamma]_\Pi + c.p. = -d^\phi(Q(\alpha, \beta, \gamma)) - ((i_{Q(\alpha, \beta, \cdot)}d^\phi\gamma) + c.p),$$

we can directly prove that

Theorem 2.7 *The triple $((A, [\cdot, \cdot], a, \phi), (A^*, [\cdot, \cdot]_\Pi, a_*, W), Q)$ is a Jacobi-quasi bialgebroid over M , which is called a triangular Jacobi-quasi bialgebroid.*

Clearly, the Lie-quasi bialgebroid associated to a twisted Poisson manifold [16] and the Jacobi-quasi bialgebroid associated to a twisted Jacobi manifold (see Theorem 2.3) are special cases of triangular Jacobi-quasi bialgebroids. Another important type of triangular quasi-Jacobi bialgebroid is the triangular quasi-Jacobi bialgebra, where Π is a solution of a Yang-Baxter's type equation.

Now, we consider the particular case where $Q = \Pi^\#(\varphi)$, with φ a d^ϕ -closed 3-form on A , and the spaces $\Gamma(A^*)$ and $\Gamma(A)$ are equipped, respectively, with the brackets

$$[\alpha, \beta]_\Pi^\varphi = [\alpha, \beta]_\Pi + \varphi(\Pi^\#(\alpha), \Pi^\#(\beta), \cdot), \quad \text{for all } \alpha, \beta \in \Gamma(A^*),$$

$[\cdot, \cdot]_\Pi$ being the Koszul bracket (5), and

$$[X, Y]' = [X, Y] - \Pi^\#(\varphi(X, Y, \cdot)), \quad \text{for all } X, Y \in \Gamma(A).$$

Under the above assumptions, by a straightforward calculation, we get:

Proposition 2.8 *The vector bundle $A^* \rightarrow M$ endowed with the structure $([\cdot, \cdot]_\Pi^\varphi, a_*)$ is a Lie algebroid over M with 1-cocycle $W = -\Pi^\#(\phi)$.*

Also, we have:

Theorem 2.9 *The triple $((A, [\cdot, \cdot]', a, \phi), (A^*, [\cdot, \cdot]_\Pi^\varphi, a_*, W), \varphi)$ is a triangular quasi-Jacobi bialgebroid over M .*

Proof. The proof is analogous to that of Theorem 8.2 in [12] and so it is omitted. $\square$

Remark 2.10 Obviously, if A is $TM \times \mathbb{R}$ equipped with the usual Lie algebroid structure with 1-cocycle, $([\cdot, \cdot], \pi, (0, 1))$, and $\Pi = (\Lambda, E) \in \Gamma(\wedge^2(TM \times \mathbb{R}))$ satisfies (4), then the manifold M is endowed with a twisted Jacobi structure. The Lie algebroid structure on $A^* = T^*M \times \mathbb{R}$ given by Proposition 2.8, is the Lie algebroid structure canonically associated with the twisted Jacobi structure on M .

3 The structure induced on the base manifold of a quasi-Jacobi bialgebroid

In this section we will investigate the structure induced on the base manifold of a quasi-Jacobi bialgebroid. Similar results hold for a Jacobi-quasi bialgebroid.

Let $((A, \phi), (A^*, W), \varphi)$ be a quasi-Jacobi bialgebroid over M . In [12], we have already considered the bracket $\{\cdot, \cdot\}$ on $C^\infty(M, \mathbb{R})$ defined, for all $f, g \in C^\infty(M, \mathbb{R})$, by

$$\{f, g\} = \langle d^\phi f, d_*^W g \rangle. \quad (6)$$

We have proved that it is $\mathbb{R}$ -bilinear, skew-symmetric and a first order differential operator on each argument [12]. On the other hand, the quasi-differential operator d on $\Gamma(\wedge A^*)$ determined by $(a, [\cdot, \cdot])$ is a derivation with respect to the usual product of functions. Therefore, the map $(f, g) \mapsto \langle df, d_* g \rangle$ is a derivation on each argument and so, there exists a bivector field Λ on M such that, for all $f, g \in C^\infty(M, \mathbb{R})$,

$$\Lambda(\delta f, \delta g) = \langle df, d_* g \rangle = -\langle dg, d_* f \rangle.$$

If E is the vector field $a_*(\phi) = -a(W)$ on M then, from (6) and because $\langle \phi, W \rangle = 0$ holds [12], we get

$$\{f, g\} = \langle d^{(0,1)} g, (\Lambda, E)^\#(d^{(0,1)} f) \rangle. \quad (7)$$

Since, for all $f \in C^\infty(M, \mathbb{R})$, $d^\phi f = (a^\phi)^*(d^{(0,1)} f)$ and $d_*^W f = (a_*^W)^*(d^{(0,1)} f)$ [12], where $(a^\phi)^*$ and $(a_*^W)^*$ denote, respectively, the transpose of a^ϕ and a_*^W , we obtain

$$(\Lambda, E)^\# = -a^\phi \circ (a_*^W)^* = a_*^W \circ (a^\phi)^*. \quad (8)$$

It is well known that any bracket of type (7) satisfies the following relation:

$$\{f, \{g, h\}\} + c.p. = \frac{1}{2}[(\Lambda, E), (\Lambda, E)]^{(0,1)}(d^{(0,1)} f, d^{(0,1)} g, d^{(0,1)} h). \quad (9)$$

Therefore, for the bracket defined by (6), in general, the Jacobi identity does not hold.

Proposition 3.1 *Let $((A, \phi), (A^*, W), \varphi)$ be a quasi-Jacobi bialgebroid over M . Then, the bracket (6) satisfies, for all $f, g, h \in C^\infty(M, \mathbb{R})$, the following identity:*

$$\{f, \{g, h\}\} + c.p. = a_*^W(\varphi)(d^{(0,1)} f, d^{(0,1)} g, d^{(0,1)} h). \quad (10)$$

In (10), a_*^W denotes the natural extension of $a_*^W : \Gamma(A^*) \rightarrow \Gamma(TM \times \mathbb{R})$ to a bundle map from $\Gamma(\wedge^3 A^*)$ to $\Gamma(\wedge^3(TM \times \mathbb{R}))$.

Proof. Let f, g and h be any three functions on $C^\infty(M, \mathbb{R})$. Taking into account the formulæ $d_*^W \{f, g\} = [d_*^W g, d_*^W f]$ (see [12]), (8) and (3), and the properties of a quasi-Jacobi bialgebroid, after a simple computation, we get

$$\begin{aligned} \{h, \{f, g\}\} &= \{\{g, f\}, h\} - \frac{1}{2}[(\Lambda, E), (\Lambda, E)]^{(0,1)}(d^{(0,1)} f, d^{(0,1)} g, d^{(0,1)} h) \\ &\quad + a_*^W(\varphi)(d^{(0,1)} f, d^{(0,1)} g, d^{(0,1)} h). \end{aligned}$$

Consequently,

$$\frac{1}{2}[(\Lambda, E), (\Lambda, E)]^{(0,1)}(d^{(0,1)} f, d^{(0,1)} g, d^{(0,1)} h) = a_*^W(\varphi)(d^{(0,1)} f, d^{(0,1)} g, d^{(0,1)} h). \quad (11)$$

Hence, from (9) and (11), we obtain (10). $\square$

Looking at equation (11), we remark that the obstruction for (M, Λ, E) to be a Jacobi manifold, i.e. to have $[(\Lambda, E), (\Lambda, E)]^{(0,1)} = (0, 0)$, is the image by a_*^W of the element φ in $\Gamma(\wedge^3 A^*)$. This obstruction can also be viewed as the image of φ under the infinitesimal action of the Lie algebroid with 1-cocycle (A^*, W) on M (see Appendix). Thus, inspired by the analogous terms of *quasi Poisson $\mathcal{G}$ -manifold* ([1], [9]) and of $(\mathcal{G}, \phi)$ -*quasi Jacobi manifold* (see Section 2.3), we say that the pair (Λ, E) defines on M a (A^*, W) -*quasi Jacobi structure*.

Thus, we have proved:

Theorem 3.2 *Let $((A, \phi), (A^*, W), \varphi)$ be a quasi-Jacobi bialgebroid over M . Then, the bracket $\{\cdot, \cdot\} : C^\infty(M, \mathbb{R}) \times C^\infty(M, \mathbb{R}) \rightarrow C^\infty(M, \mathbb{R})$ given by*

$$\{f, g\} = \langle d^\phi f, d_*^W g \rangle, \quad \text{for } f, g \in C^\infty(M, \mathbb{R}),$$

defines a (A^, W) -quasi Jacobi structure on M .*

Remark 3.3 In the case where $((A, \phi), (A^*, W), Q)$ is a Jacobi-quasi bialgebroid over M , we can easily prove that the Jacobi identity of the bracket defined by (6) is violated by the image of Q under a^ϕ . For this reason, we shall call the structure (Λ, E) induced on M , an (A, ϕ) -*quasi Jacobi structure*. We note that, for the proof of this result, we use the relation $[d^\phi f, d^\phi g]_* = d^\phi \{f, g\}$, $f, g \in C^\infty(M, \mathbb{R})$, which leads to

$$(\Lambda, E)^\# = a^\phi \circ (a_*^W)^* = -a_*^W \circ (a^\phi)^*. \quad (12)$$

Examples 3.4

1) *A^* -Quasi Poisson structures:* If $((A, \phi), (A^*, W), \varphi)$ is a quasi-Lie bialgebroid over M , i.e. both 1-cocycles ϕ and W are zero, Theorem 3.2 establishes the existence of a structure on M , defined by the bracket

$$\{f, g\} = \langle df, d_* g \rangle, \quad f, g \in C^\infty(M, \mathbb{R}),$$

on $C^\infty(M, \mathbb{R})$, which is associated to a bivector field Λ on M satisfying $[\Lambda, \Lambda] = 2a_*(\varphi)$. In our terminology, Λ endows M with a A^* -*quasi Poisson structure*. We remark that this result was obtained in [6] by different techniques.

2) *Jacobi structures:* When $\varphi = 0$, i.e. $((A, \phi), (A^*, W))$ is a Jacobi bialgebroid over M , the structure (Λ, E) on M determined by Theorem 3.2 is a Jacobi structure, and we recover the well known result of [5].

3) *Twisted Jacobi structures:* When φ is the image of an element $(\varphi_M, \omega_M) \in \Gamma(\wedge^3(T^*M \times \mathbb{R}))$ by the transpose map $(a^\phi)^* : \Gamma(\wedge^3(T^*M \times \mathbb{R})) \rightarrow \Gamma(\wedge^3 A^*)$ of a^ϕ , i.e. $\varphi = (a^\phi)^*(\varphi_M, \omega_M)$, then,

$$\frac{1}{2}[(\Lambda, E), (\Lambda, E)]^{(0,1)} = a_*^W(\varphi) = a_*^W((a^\phi)^*(\varphi_M, \omega_M)) \stackrel{(8)}{=} (\Lambda, E)^\#(\varphi_M, \omega_M).$$

Also, we have

$$\underbrace{d^\phi((a^\phi)^*(\varphi_M, \omega_M))}_{=\varphi} = 0 \Leftrightarrow (a^\phi)^*(d^{(0,1)}(\varphi_M, \omega_M)) = 0,$$

which means that (φ_M, ω_M) is $d^{(0,1)}$ -closed on the distribution $Im(a^\phi)$. This distribution is not, in general, involutive due to condition 2) of Definition 2.1. However, when $Im(a^\phi)$ is involutive, as in the case where a^ϕ is surjective, $((\Lambda, E), \omega_M)$ defines a twisted Jacobi structure on the leaves of $Im(a^\phi)$.

4) *The case of the quasi-Jacobi bialgebroid associated to a twisted Jacobi manifold:* Let $(M, (\Lambda_1, E_1), \omega)$ be a twisted Jacobi manifold and let $((TM \times \mathbb{R}, [\cdot, \cdot]', \pi, (0, 1)), (T^*M \times \mathbb{R}, [\cdot, \cdot]_{(\Lambda_1, E_1)}^\omega, \pi \circ (\Lambda_1, E_1)^\#, (-E_1, 0)), (\delta\omega, \omega))$ be its associated quasi-Jacobi bialgebroid. Then, the $(T^*M \times \mathbb{R}, (-E_1, 0))$ -quasi Jacobi structure induced on M coincides with the initial one (Λ_1, E_1) . In fact, for any $f, g \in C^\infty(M, \mathbb{R})$ and taking into account that $d^{(0,1)}f = d^{(0,1)}f$, d' being the quasi-differential of $TM \times \mathbb{R}$ determined by the structure $([\cdot, \cdot]', \pi)$, and that $(d_*^\omega)^{(-E_1, 0)}g = -(\Lambda_1, E_1)^\#(d^{(0,1)}g)$, we have

$$\{f, g\} = \langle d^{(0,1)}f, (d_*^\omega)^{(-E_1, 0)}g \rangle = \langle d^{(0,1)}f, -(\Lambda_1, E_1)^\#(d^{(0,1)}g) \rangle = \{f, g\}_1,$$

where $\{\cdot, \cdot\}_1$ denotes the bracket associated to (Λ_1, E_1) .

Moreover, if we consider the Jacobi-quasi bialgebroid $((TM \times \mathbb{R}, [\cdot, \cdot]', \pi, (0, 1)), (T^*M \times \mathbb{R}, [\cdot, \cdot]_{(\Lambda_1, E_1)}^\omega, \pi \circ (\Lambda_1, E_1)^\#, (-E_1, 0)), (\Lambda_1, E_1)^\#(\delta\omega, \omega))$ associated to the twisted Jacobi manifold $(M, (\Lambda_1, E_1), \omega)$, we get that the $(TM \times \mathbb{R}, (0, 1))$ -quasi Jacobi structure (Λ, E) induced on M is the opposite of (Λ_1, E_1) . It suffices to remark that

$$\begin{aligned} (\Lambda, E)^\# &\stackrel{(12)}{=} \pi^{(0,1)} \circ ((\pi \circ (\Lambda_1, E_1)^\#)^{(-E_1, 0)})^* \\ &= \pi^{(0,1)} \circ ((\Lambda_1, E_1)^\#)^* \circ (\pi^{(0,1)})^* = -(\Lambda_1, E_1)^\#. \end{aligned} \quad (13)$$

5) *The induced structure on a quasi Jacobi manifold:* We consider the Jacobi-quasi bialgebroid $((TM \times \mathbb{R}, [\cdot, \cdot]', \pi, (0, 1)), (T^*M \times \mathbb{R}, [\cdot, \cdot]_{(\Lambda_1, E_1)}^\omega, \pi \circ (\Lambda_1, E_1)^\#, (-E_1, 0)), a^\phi(Q_\psi))$ associated to a $(\mathcal{G}, \phi)$ -quasi Jacobi manifold (M, Λ_1, E_1) . Then, repeating the computation (13), we conclude that, as in the previous case, the $(TM \times \mathbb{R}, (0, 1))$ -quasi Jacobi structure (Λ, E) induced on M is the opposite of (Λ_1, E_1) .

6) *The case of a triangular quasi-Jacobi bialgebroid:* If we consider a triangular quasi-Jacobi bialgebroid over M of type $((A, [\cdot, \cdot]', a, \phi), (A^*, [\cdot, \cdot]_\Pi^\varphi, a_*, W), \varphi)$, presented in Theorem 2.9, then, for all $f \in C^\infty(M, \mathbb{R})$,

$$d_*^W f = -\Pi^\#(d^\phi f) = -(\Pi^\# \circ (a^\phi)^*)(d^{(0,1)}f).$$

So, the bracket (6) in $C^\infty(M, \mathbb{R})$ is given by

$$\{f, g\} = \langle d^\phi f, d_*^W g \rangle = \langle d^{(0,1)}g, (a^\phi \circ \Pi^\# \circ (a^\phi)^*)d^{(0,1)}f \rangle.$$

On the other hand, considering the (A^*, W) -quasi Jacobi structure (Λ, E) on M , we also have

$$\{f, g\} = \langle d^{(0,1)}g, (\Lambda, E)^\#(d^{(0,1)}f) \rangle.$$

Hence,

$$(\Lambda, E)^\# = a^\phi \circ \Pi^\# \circ (a^\phi)^*,$$

which means that (Λ, E) is the image by a^ϕ of Π and that a^ϕ is a type of “twisted Jacobi morphism” between (A, ϕ, Π) and $(TM \times \mathbb{R}, (0, 1), (\Lambda, E))$.

4 Quasi-Lie bialgebroids associated to quasi-Jacobi bialgebroids

Given a Lie algebroid $(A, [\cdot, \cdot], a)$ over M , we can endow the vector bundle $\tilde{A} = A \times \mathbb{R} \rightarrow M \times \mathbb{R}$ with a Lie algebroid structure over $M \times \mathbb{R}$ as follows. The sections of $\tilde{A}$ can be identified with the t -dependent sections of A , t being the canonical coordinate on $\mathbb{R}$, i.e., for any $\tilde{X} \in \Gamma(\tilde{A})$ and $(x, t) \in M \times \mathbb{R}$, $\tilde{X}(x, t) = \tilde{X}_t(x)$, where $\tilde{X}_t \in \Gamma(A)$. This identification induces, in a natural way, a Lie bracket on $\Gamma(\tilde{A})$, also denoted by $[\cdot, \cdot]$:

$$[\tilde{X}, \tilde{Y}](x, t) = [\tilde{X}_t, \tilde{Y}_t](x), \quad \tilde{X}, \tilde{Y} \in \Gamma(\tilde{A}), \quad (x, t) \in M \times \mathbb{R},$$

and a bundle map, also denoted by a , $a : \tilde{A} \rightarrow T(M \times \mathbb{R}) \equiv TM \oplus T\mathbb{R}$ with $a(\tilde{X}) = a(\tilde{X}_t)$, in such a way that $(\tilde{A}, [\cdot, \cdot], a)$ becomes a Lie algebroid over $M \times \mathbb{R}$. If ϕ is a 1-cocycle of the Lie algebroid A , we know from [5] that $\tilde{A}$ can be equipped with two other Lie algebroid structures over $M \times \mathbb{R}$, $([\cdot, \cdot]^\phi, \tilde{a}^\phi)$ and $([\cdot, \cdot]^\phi, \hat{a}^\phi)$ given, for all $\tilde{X}, \tilde{Y} \in \Gamma(\tilde{A})$, by

$$[\tilde{X}, \tilde{Y}]^\phi = [\tilde{X}_t, \tilde{Y}_t] + \langle \phi, \tilde{X}_t \rangle \frac{\partial \tilde{Y}}{\partial t} - \langle \phi, \tilde{Y}_t \rangle \frac{\partial \tilde{X}}{\partial t}; \quad (14)$$

$$\tilde{a}^\phi(\tilde{X}) = a(\tilde{X}_t) + \langle \phi, \tilde{X}_t \rangle \frac{\partial}{\partial t}; \quad (15)$$

and

$$[\tilde{X}, \tilde{Y}]^{\hat{\phi}} = e^{-t} \left([\tilde{X}_t, \tilde{Y}_t] + \langle \phi, \tilde{X}_t \rangle \left(\frac{\partial \tilde{Y}}{\partial t} - \tilde{Y} \right) - \langle \phi, \tilde{Y}_t \rangle \left(\frac{\partial \tilde{X}}{\partial t} - \tilde{X} \right) \right); \quad (16)$$

$$\hat{a}^\phi(\tilde{X}) = e^{-t}(\tilde{a}^\phi(\tilde{X})). \quad (17)$$

Let $((A, \phi), (A^*, W), \varphi)$ be a quasi-Jacobi bialgebroid over M . Then, $(A^*, [\cdot, \cdot]_*, a_*, W)$ is a Lie algebroid with 1-cocycle and we can consider on $\tilde{A}^*$ the Lie algebroid structure $([\cdot, \cdot]_*^W, \hat{a}_*^W)$ defined by (16) and (17). Although A is not endowed with a Lie algebroid structure, we can still consider on $\Gamma(\tilde{A})$ a bracket $[\cdot, \cdot]^\phi$ and a bundle map $\tilde{a}^\phi$ given by (14) and (15), respectively. We set $\tilde{\varphi} = e^t \varphi$.

Theorem 4.1 *Under the above assumptions, we have:*

- 1) *The triple $((A, \phi), (A^*, W), \varphi)$ is a quasi-Jacobi bialgebroid over M if and only if $(\tilde{A}, \tilde{A}^*, \tilde{\varphi})$ is a quasi-Lie bialgebroid over $M \times \mathbb{R}$.*
- 2) *If $\tilde{\Lambda}$ is the induced $\tilde{A}^*$ -quasi Poisson structure on $M \times \mathbb{R}$, then it is the “quasi Poissonization” of the induced (A^*, W) -quasi Jacobi structure (Λ, E) on M .*

Proof. 1) Let us suppose that $((A, \phi), (A^*, W), \varphi)$ is a quasi-Jacobi bialgebroid over M and let $\tilde{X}, \tilde{Y}$ and $\tilde{Z}$ be three arbitrary sections in $\Gamma(\tilde{A})$ and $\tilde{f} \in C^\infty(M \times \mathbb{R}, \mathbb{R})$. A straightforward computation gives

$$[\tilde{X}, \tilde{f}\tilde{Y}]^\phi = \tilde{f}[\tilde{X}, \tilde{Y}]^\phi + (\tilde{a}^\phi(\tilde{X})\tilde{f})\tilde{Y}. \quad (18)$$

Moreover,

$$\begin{aligned}
\tilde{a}^\phi([\tilde{X}, \tilde{Y}]^{\sim\phi}) &= [a(\tilde{X}_t), a(\tilde{Y}_t)] - a_*(\varphi(\tilde{X}_t, \tilde{Y}_t, \cdot)) + \langle \phi, [\tilde{X}_t, \tilde{Y}_t] \rangle \frac{\partial}{\partial t} \\
&\quad + \langle \phi, \tilde{X} \rangle \left(a\left(\frac{\partial \tilde{Y}}{\partial t}\right) + \langle \phi, \frac{\partial \tilde{Y}}{\partial t} \rangle \frac{\partial}{\partial t} \right) - \langle \phi, \tilde{Y} \rangle \left(a\left(\frac{\partial \tilde{X}}{\partial t}\right) + \langle \phi, \frac{\partial \tilde{X}}{\partial t} \rangle \frac{\partial}{\partial t} \right) \\
&= [\tilde{a}^\phi(\tilde{X}), \tilde{a}^\phi(\tilde{Y})] - \hat{a}_*^W(\tilde{\varphi}(\tilde{X}, \tilde{Y}, \cdot)),
\end{aligned}$$

where, in the last equality, we have used $d\phi - \varphi(W, \cdot, \cdot) = 0$. On the other hand,

$$\begin{aligned}
[[\tilde{X}, \tilde{Y}]^{\sim\phi}, \tilde{Z}]^{\sim\phi} + c.p. &= \left([[\tilde{X}_t, \tilde{Y}_t], \tilde{Z}_t] - d\phi(\tilde{X}_t, \tilde{Y}_t) \frac{\partial \tilde{Z}_t}{\partial t} \right) + c.p. \\
&= -d_*^W(\varphi(\tilde{X}_t, \tilde{Y}_t, \tilde{Z}_t)) - \left(\left(i_{\varphi(\tilde{X}_t, \tilde{Y}_t, \cdot)} d_*^W \tilde{Z}_t + \varphi(W, \tilde{X}_t, \tilde{Y}_t) \frac{\partial \tilde{Z}_t}{\partial t} \right) + c.p. \right) \\
&= -\hat{d}_*^W(\tilde{\varphi}(\tilde{X}, \tilde{Y}, \tilde{Z})) - \left(i_{\tilde{\varphi}(\tilde{X}, \tilde{Y}, \cdot)} \hat{d}_*^W \tilde{Z} + c.p. \right), \tag{19}
\end{aligned}$$

where $\hat{d}_*^W$ is the differential operator of $(\tilde{A}^*, [\cdot, \cdot]_*^W, \hat{a}_*^W)$. Because $d^\phi\varphi = 0$, we get

$$\tilde{d}^\phi \tilde{\varphi} = 0, \tag{20}$$

where $\tilde{d}^\phi$ is the quasi-differential operator determined by the structure $([\cdot, \cdot]_*^{\sim\phi}, \tilde{a}^\phi)$ on $\tilde{A}$. Finally, after a very long computation we obtain

$$\hat{d}_*^W[\tilde{P}, \tilde{Q}]^{\sim\phi} = [\hat{d}_*^W \tilde{P}, \tilde{Q}]^{\sim\phi} + (-1)^{p+1}[\tilde{P}, \hat{d}_*^W \tilde{Q}]^{\sim\phi}, \tag{21}$$

for $\tilde{P} \in \Gamma(\bigwedge^p \tilde{A})$ and $\tilde{Q} \in \Gamma(\bigwedge \tilde{A})$. From relations (18) to (21), we conclude that $(\tilde{A}, \tilde{A}^*, \tilde{\varphi})$ is a quasi-Lie bialgebroid over $M \times \mathbb{R}$.

Now, let us suppose that $(\tilde{A}, \tilde{A}^*, \tilde{\varphi})$ is a quasi-Lie bialgebroid over $M \times \mathbb{R}$ and take three sections X, Y and Z of A and $f \in C^\infty(M, \mathbb{R})$. These sections can be viewed as sections of $\tilde{A}$ that don't depend on t , as well as the function f can also be viewed as a function on $C^\infty(M \times \mathbb{R}, \mathbb{R})$. Condition 1) of Definition 2.1 is immediate from $[X, fY]^{\sim\phi} = f[X, Y]^{\sim\phi} + (\tilde{a}^\phi(X)f)Y$. The condition $\tilde{a}^\phi([X, Y]^{\sim\phi}) = [\tilde{a}^\phi(X), \tilde{a}^\phi(Y)] - \hat{a}_*^W(\tilde{\varphi}(X, Y, \cdot))$ is equivalent to conditions 2) and 4) of Definition 2.1. From $\tilde{d}^\phi \tilde{\varphi} = 0$ we deduce $d^\phi\varphi = 0$. Finally, by similar computations, we obtain the two remaining conditions that lead to the conclusion that $((A, \phi), (A^*, W), \varphi)$ is a quasi-Jacobi bialgebroid over M .

2) Let $\tilde{\Lambda}$ be the $\tilde{A}^*$ -quasi Poisson structure induced by $(\tilde{A}, \tilde{A}^*, \tilde{\varphi})$ on $M \times \mathbb{R}$. For all $\tilde{f}, \tilde{g} \in C^\infty(M \times \mathbb{R}, \mathbb{R})$, we have

$$\{\tilde{f}, \tilde{g}\} = \tilde{\Lambda}(\delta\tilde{f}, \delta\tilde{g})$$

and, on the other hand,

$$\{\tilde{f}, \tilde{g}\} = \langle \tilde{d}^\phi \tilde{f}, \hat{d}_*^W \tilde{g} \rangle = e^{-t}(\langle d\tilde{f}, d_*\tilde{g} \rangle + \frac{\partial \tilde{f}}{\partial t} a_*(\phi)(\tilde{g}) + \frac{\partial \tilde{g}}{\partial t} a(W)(\tilde{f})).$$

If (Λ, E) is the (A^*, W) -quasi Jacobi structure induced by $((A, \phi), (A^*, W), \varphi)$ on M , since $E = a_*(\phi) = -a(W)$ and $\Lambda(\delta\tilde{f}, \delta\tilde{g}) = \langle d\tilde{f}, d_*\tilde{g} \rangle$, we get that $\tilde{\Lambda} = e^{-t}(\Lambda + \frac{\partial}{\partial t} \wedge E)$. $\square$

For the case of Jacobi-quasi bialgebroids we can prove a similar result. Let $((A, \phi), (A^*, W), Q)$ be a Jacobi-quasi bialgebroid over M . We consider on $\tilde{A}$ the Lie algebroid structure $([\cdot, \cdot]^\phi, \hat{a}^\phi)$ defined by (16) and (17), on $\tilde{A}^*$ the structure $([\cdot, \cdot]_*^W, \tilde{a}_*^W)$ defined by (14) and (15), and we set $\tilde{Q} = e^t Q$.

Theorem 4.2 *Under the above assumptions, we have:*

- 1) *The triple $((A, \phi), (A^*, W), Q)$ is a Jacobi-quasi bialgebroid over M if and only if $(\tilde{A}, \tilde{A}^*, \tilde{Q})$ is a Lie-quasi bialgebroid over $M \times \mathbb{R}$.*
- 2) *If $\tilde{\Lambda}$ is the $\tilde{A}$ -quasi Poisson structure induced on $M \times \mathbb{R}$, then it is the “quasi Poissonization” of the (A, ϕ) -quasi Jacobi structure (Λ, E) induced on M .*

5 Appendix: Action of a Lie algebroid with 1-cocycle

In this Appendix, we extend the definition of Lie algebroid action [10] to that of Lie algebroid with 1-cocycle action.

Let $(A, [\cdot, \cdot], a)$ be a Lie algebroid on M and $\varpi : F \rightarrow M$ a fibered manifold with base M , i.e. $\varpi : F \rightarrow M$ is a surjective submersion onto M . We recall that an *infinitesimal action of A on F* ([10]) is a $\mathbb{R}$ -linear map $\mathbf{ac} : \Gamma(A) \rightarrow \Gamma(TF)$ such that:

- (1) for each $X \in \Gamma(A)$, $\mathbf{ac}(X)$ is projectable to $a(X)$,
- (2) the map $\mathbf{ac}$ preserves brackets,
- (3) $\mathbf{ac}(fX) = (f \circ \varpi)\mathbf{ac}(X)$, for all $f \in C^\infty(M, \mathbb{R})$ and $X \in \Gamma(A)$.

Definition 5.1 *An infinitesimal action of (A, ϕ) on F is a $\mathbb{R}$ -linear map $\mathbf{ac}^\phi : \Gamma(A) \rightarrow \Gamma(TF \times \mathbb{R})$ given, for each $X \in \Gamma(A)$, by*

$$\mathbf{ac}^\phi(X) = \mathbf{ac}(X) + \langle \phi, X \rangle,$$

where $\mathbf{ac}$ is an infinitesimal action of A on F .

In the particular case where M is a point and therefore A is a Lie algebra, we obtain, from Definition 5.1, the notion of infinitesimal action of a Lie algebra with 1-cocycle on a manifold F , used in the definition of quasi Jacobi structures, in section 2.3.

If, in the Definition 5.1, $F = M$ and $\varpi : M \rightarrow M$ is the identity, we get the concept of infinitesimal action of (A, ϕ) on the base manifold M that we have used to characterize the structure induced on the base manifold of a quasi-Jacobi bialgebroid.

Acknowledgments Research of J. M. Nunes da Costa supported by CMUC-FCT and POCI/MAT/58452/2004.

References

- [1] Alekseev, A., Kosmann-Schwarzbach, Y., Meinrenken, E.: Quasi-Poisson manifolds. *Canad. J. Math.* **54**, 3-29 (2002)

- [2] Drinfeld, V.G.: Quasi-Hopf algebras. Leningrad Math. J. **1**, 1419-1457 (1990)
- [3] Grabowski, J., Marmo, G.: Jacobi structures revisited. J. Phys. A : Math. Gen. **34**, 10975-10990 (2001)
- [4] Grabowski, J., Marmo, G.: The graded Jacobi algebras and (co)homology. J. Phys. A : Math. Gen. **36**, 161-181 (2003)
- [5] Iglesias, D., Marrero, J.C.: Generalized Lie bialgebroids and Jacobi structures. J. Geom. Phys. **40**, 176-200 (2001)
- [6] Iglesias Ponte, D., Laurent-Gengoux, C., Xu, P.: Universal lifting theorem and quasi-Poisson groupoids. ArXiv: math.DG/0507396
- [7] Kerbrat, Y., Souici-Benhamadi, Z.: Variétés de Jacobi et groupoïdes de contact. C. R. Acad. Sci. Paris, Série I, **317**, 81-86 (1993)
- [8] Klimčík, C., Strobl, T.: WZW-Poisson manifolds. J. Geom. Phys. **43**, 341 (2002)
- [9] Kosmann-Schwarzbach, Y.: Quasi, twisted, and all that ... In Poisson geometry and Lie algebroid theory. In: Marsden, J.E., Ratiu, T.S. (eds.) The breadth of symplectic and Poisson geometry. Progress in Mathematics, vol. 232, pp. 363-389. Birkhauser, Boston (2005)
- [10] Kosmann-Schwarzbach, Y., Mackenzie, K.: Differential operators and actions of Lie algebroids. Contemp. Math. **315**, 213-233 (2002)
- [11] Nunes da Costa, J.M. and Clemente-Gallardo, J.: Dirac structures for generalized Lie bialgebroids. J. Phys. A : Math. Gen. **37**, 2671-2692 (2004)
- [12] Nunes da Costa, J.M., Petalidou, F.: Twisted Jacobi manifolds, twisted Dirac-Jacobi structures and quasi-Jacobi bialgebroids. J. Phys. A: Math. Gen. **39**, 10449-10475 (2006)
- [13] Nunes da Costa, J.M., Petalidou, F.: Characteristic foliation of twisted Jacobi manifolds. ArXiv: math.DG/0612145.
- [14] Park, J.S.: Topological open p -branes. In: Fukaya, K., Oh, Y.G., Ono, K., Tian, G. (eds.) Symplectic geometry and Mirror Symmetry (Seoul, 2000). World Sci. Publishing, River Edge, NJ, pp. 311-384 (2001)
- [15] Roytenberg, D.: Courant algebroids, derived brackets and even symplectic supermanifolds. Ph.D. Thesis, University of California, Berkeley (1999) (arXiv: math.DG/9910078).
- [16] Roytenberg, D.: Quasi-Lie Bialgebroids and Twisted Poisson Manifolds. Lett. Math. Phys. **61**, 123-137 (2002)
- [17] Ševera, P., Weinstein, A.: Poisson geometry with a 3-form background. In: Noncommutative geometry and string theory. Prog. Theor. Phys., Suppl. **144**, 145-154 (2001)